


Amendments to the Claims:

This listing of claims will replace all prior versions and listings of claims in the application:

Listing of Claims:

Claims 1 to 17 (canceled)

Claim 18 (new): A compound of Formula (I), or a salt, pro-drug, or solvate thereof:



wherein:

R¹ is methoxymethyl;

R² is selected from -C(O)NR⁴R⁵, -SO₂NR⁴R⁵, and -S(O)_pR⁴;

HET-1 is a 5- or 6-membered, C-linked heteroaryl ring containing a nitrogen atom in the 2-position and optionally 1 or 2 further ring heteroatoms independently selected from O, N, and S; which ring is optionally substituted on an available carbon atom, or on a ring nitrogen atom provided it is not thereby quaternised, with 1 or 2 substituents independently selected from R⁶;

R³ is selected from halo, fluoromethyl, difluoromethyl, trifluoromethyl, methyl, methoxy, and cyano;

R⁴ is selected from hydrogen and (1-4C)alkyl;

R⁵ is hydrogen or (1-4C)alkyl;

R⁶ is independently selected from (1-4C)alkyl, halo, hydroxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkylS(O)_p(1-4C)alkyl, amino(1-4C)alkyl, (1-4C)alkylamino(1-4C)alkyl, di(1-4C)alkylamino(1-4C)alkyl, and HET-4;

HET-4 is a 5- or 6-membered, C- or N- linked unsubstituted heteroaryl ring containing 1, 2, or 3

ring heteroatoms independently selected from O, N, and S;

p is independently at each occurrence 0, 1, or 2;

m is 0 or 1;

n is 0, 1, or 2;

provided that when **m** is 0, then **n** is 1 or 2.

Claim 19 (new): A compound of Formula (I), as claimed in Claim 18, which is selected from:

- 3-{2-chloro-4-[(dimethylamino)sulfonyl]phenoxy}-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-[(2-chloro-4-[(1-methylethyl)amino]sulfonyl}phenyl)oxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-{4-[(dimethylamino)sulfonyl]phenoxy}-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-{4-[(1-methylethyl)amino]sulfonyl}phenoxy}-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-(4-cyanophenoxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-{[4-(aminocarbonyl)phenyl]oxy}-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-[4-(ethylsulfonyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide;
- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-{[3-(methylthio)phenyl]oxy}benzamide;
- 3-({4-[(1-methylethyl)thio]phenyl}oxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-[3-(methylsulfonyl)phenoxy]benzamide;
- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-[3-(methylsulfinyl)phenoxy]benzamide;
- 3-({4-[(1-methylethyl)sulfonyl]phenyl}oxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-

methyl-1H-pyrazol-3-yl)benzamide;

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-[4-(methylsulfonyl)phenoxy]benzamide;

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]-N-1,3-thiazol-2-ylbenzamide;

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]-N-(4-methyl-1,3-thiazol-2-yl)benzamide;

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]-N-(5-methyl-1,3-thiazol-2-yl)benzamide;

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]-N-(5-methyl-1,3,4-thiadiazol-2-yl)benzamide;

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]-N-(3-methyl-1,2,4-thiadiazol-5-yl)benzamide;

N-(1-ethyl-1H-pyrazol-3-yl)-3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]benzamide;

3-(3,5-difluorophenoxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;

N-(5-bromopyridin-2-yl)-3-(3,5-difluorophenoxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]benzamide;

3-(3,5-difluorophenoxy)-N-[4-(hydroxymethyl)-1,3-thiazol-2-yl]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]benzamide;

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]-N-1H-pyrazol-3-ylbenzamide;

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(5-methyl-1H-pyrazol-3-yl)-5-[4-(methylsulfonyl)phenoxy]benzamide;

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-[4-(methoxymethyl)-1,3-thiazol-2-yl]-5-[4-(methylsulfonyl)phenoxy]benzamide;

3-{4-[(dimethylamino)carbonyl]phenoxy}-5-[(1S)-2-methoxy-1-methylethoxy]-N-(3-methyl-1,2,4-thiadiazol-5-yl)benzamide;

3-{4-[(dimethylamino)carbonyl]phenoxy}-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;

2-methoxy-4-(3-[(1S)-2-methoxy-1-methylethoxy]-5-{[(1-methyl-1H-pyrazol-3-yl)amino]carbonyl}phenoxy)-N-methylbenzamide;

2-methoxy-4-(3-[(1S)-2-methoxy-1-methylethoxy]-5-{[(1-methyl-1H-pyrazol-3-yl)amino]carbonyl}phenoxy)-N,N-dimethylbenzamide;

3-{4-[(dimethylamino)carbonyl]phenoxy}-5-[(1S)-2-methoxy-1-methylethoxy]-N-1H-pyrazol-3-ylbenzamide;

3-fluoro-4-{3-[(1S)-2-methoxy-1-methylethoxy]-5-[(1H-pyrazol-3-ylamino)carbonyl]phenoxy}-N,N-dimethylbenzamide;

3-{4-[(dimethylamino)carbonyl]phenoxy}-5-[(1S)-2-methoxy-1-methylethoxy]-N-(5-methyl-1H-pyrazol-3-yl)benzamide;

3-[4-(ethylsulfonyl)-2-fluorophenoxy]-5-[(1S)-2-methoxy-1-methylethoxy]-N-1H-pyrazol-3-ylbenzamide;

3-[2-fluoro-4-(methylsulfonyl)phenoxy]-5-[(1S)-2-methoxy-1-methylethoxy]-N-1H-pyrazol-3-ylbenzamide; and

3-[4-(ethylsulfonyl)-2-fluorophenoxy]-5-[(1S)-2-methoxy-1-methylethoxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;

or a salt, pro-drug, or solvate thereof.

Claim 20 (new): A compound of Formula (I), as claimed in Claim 19, which is selected from:

3-{2-chloro-4-[(dimethylamino)sulfonyl]phenoxy}-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;

3-[(2-chloro-4-{[(1-methylethyl)amino]sulfonyl}phenyl)oxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;

3-{4-[(dimethylamino)sulfonyl]phenoxy}-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;

3-{4-[(1-methylethyl)amino]sulfonyl}phenoxy}-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;

3-(4-cyanophenoxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;

3-{[4-(aminocarbonyl)phenyl]oxy}-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-

pyrazol-3-yl)benzamide;

3-[4-(ethylsulfonyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide;

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-{[3-(methylthio)phenyl]oxy}benzamide;

3-({4-[(1-methylethyl)thio]phenyl}oxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-[3-(methylsulfonyl)phenoxy]benzamide;

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-[3-(methylsulfinyl)phenoxy]benzamide;

3-({4-[(1-methylethyl)sulfonyl]phenyl}oxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-[4-(methylsulfonyl)phenoxy]benzamide;

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]-N-1,3-thiazol-2-ylbenzamide;

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]-N-(4-methyl-1,3-thiazol-2-yl)benzamide;

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]-N-(5-methyl-1,3-thiazol-2-yl)benzamide;

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]-N-(5-methyl-1,3,4-thiadiazol-2-yl)benzamide;

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]-N-(3-methyl-1,2,4-thiadiazol-5-yl)benzamide;

N-(1-ethyl-1H-pyrazol-3-yl)-3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]benzamide;

3-(3,5-difluorophenoxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;

N-(5-bromopyridin-2-yl)-3-(3,5-difluorophenoxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]benzamide;

3-(3,5-difluorophenoxy)-N-[4-(hydroxymethyl)-1,3-thiazol-2-yl]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]benzamide;

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(5-methyl-1H-pyrazol-3-yl)-5-[4-(methylsulfonyl)phenoxy]benzamide;

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-[4-(methoxymethyl)-1,3-thiazol-2-yl]-5-[4-(methylsulfonyl)phenoxy]benzamide;

3-{4-[(dimethylamino)carbonyl]phenoxy}-5-[(1S)-2-methoxy-1-methylethoxy]-N-(3-methyl-1,2,4-thiadiazol-5-yl)benzamide;

3-{4-[(dimethylamino)carbonyl]phenoxy}-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;

2-methoxy-4-(3-[(1S)-2-methoxy-1-methylethoxy]-5-[(1-methyl-1H-pyrazol-3-yl)amino]carbonyl}phenoxy)-N-methylbenzamide;

2-methoxy-4-(3-[(1S)-2-methoxy-1-methylethoxy]-5-[(1-methyl-1H-pyrazol-3-yl)amino]carbonyl}phenoxy)-N,N-dimethylbenzamide;

3-{4-[(dimethylamino)carbonyl]phenoxy}-5-[(1S)-2-methoxy-1-methylethoxy]-N-1H-pyrazol-3-ylbenzamide;

3-fluoro-4-{3-[(1S)-2-methoxy-1-methylethoxy]-5-[(1H-pyrazol-3-ylamino)carbonyl]phenoxy}-N,N-dimethylbenzamide;

3-{4-[(dimethylamino)carbonyl]phenoxy}-5-[(1S)-2-methoxy-1-methylethoxy]-N-(5-methyl-1H-pyrazol-3-yl)benzamide;

3-[4-(ethylsulfonyl)-2-fluorophenoxy]-5-[(1S)-2-methoxy-1-methylethoxy]-N-1H-pyrazol-3-ylbenzamide;

3-[2-fluoro-4-(methylsulfonyl)phenoxy]-5-[(1S)-2-methoxy-1-methylethoxy]-N-1H-pyrazol-3-ylbenzamide; and

3-[4-(ethylsulfonyl)-2-fluorophenoxy]-5-[(1S)-2-methoxy-1-methylethoxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;


or a salt, pro-drug, or solvate thereof.

Claim 21 (new): A compound of Formula (I) as claimed in Claim 18, which is:

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]-N-(3-methyl-1,2,4-thiadiazol-5-yl)benzamide;

or a salt, pro-drug, or solvate thereof.

Claim 22 (new): A compound of Formula (I) or a salt, pro-drug, or solvate thereof:



wherein:

R¹ is methoxymethyl;

R² is selected from -C(O)-HET-3 and -SO₂-HET-3;

HET-1 is a 5- or 6-membered, C-linked heteroaryl ring containing a nitrogen atom in the 2-position and optionally 1 or 2 further ring heteroatoms independently selected from O, N, and S; which ring is optionally substituted on an available carbon atom, or on a ring nitrogen atom provided it is not thereby quaternised, with 1 or 2 substituents independently selected from R⁶;

HET-2 is a 4-, 5-, or 6-membered, C- or N-linked heterocyclyl ring containing 1, 2, 3, or 4 heteroatoms independently selected from O, N and S, wherein a -CH₂- group can optionally be replaced by a -C(O)-, and wherein a sulphur atom in the heterocyclic ring may optionally be oxidised to a S(O) or S(O)₂ group, which ring is optionally substituted on an available carbon or nitrogen atom by 1 or 2 substituents independently selected from R⁷;

R³ is selected from halo, fluoromethyl, difluoromethyl, trifluoromethyl, methyl, methoxy, and cyano;

R⁴ is selected from hydrogen; (1-4C)alkyl optionally substituted with 1 or 2 substituents independently selected from HET-2, -OR⁵, -SO₂R⁵, (3-6C)cycloalkyl (optionally substituted with 1 group selected from R⁷), and -C(O)NR⁵R⁵; (3-6C)cycloalkyl (optionally substituted with 1 group selected from R⁷); and HET-2;

R⁵ is hydrogen or (1-4C)alkyl;

or **R⁴** and **R⁵** together with the nitrogen atom to which they are attached may form a heterocyclyl

ring system as defined by HET-3;

R⁶ is independently selected from (1-4C)alkyl, halo, hydroxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkylS(O)_p(1-4C)alkyl, amino(1-4C)alkyl, (1-4C)alkylamino(1-4C)alkyl, di(1-4C)alkylamino(1-4C)alkyl, and HET-4;

R⁷ is selected from -OR⁵, (1-4C)alkyl, -C(O)(1-4C)alkyl, -C(O)NR⁴R⁵, (1-4C)alkoxy(1-4C)alkyl, hydroxy(1-4C)alkyl, and -S(O)_pR⁵;

HET-3 is an N-linked, 4-, 5-, or 6-membered, saturated or partially unsaturated heterocyclyl ring, optionally containing 1 or 2 further heteroatoms independently selected from O, N, and S, wherein a -CH₂- group can optionally be replaced by a -C(O)- and wherein a sulphur atom in the ring may optionally be oxidised to a S(O) or S(O)₂ group; which ring is optionally substituted on an available carbon or nitrogen atom by 1 or 2 substituents independently selected from R⁸; or

HET-3 is an N-linked, 7-membered, saturated or partially unsaturated heterocyclyl ring, optionally containing 1 further heteroatom independently selected from O, S, and N, wherein a -CH₂- group can optionally be replaced by a -C(O)- group and wherein a sulphur atom in the ring may optionally be oxidised to a S(O) or S(O)₂ group; which ring is optionally substituted on an available carbon or nitrogen atom by 1 or 2 substituents independently selected from R⁸; or

HET-3 is an N-linked, 6- to 10-membered bicyclic saturated or partially unsaturated heterocyclyl ring, optionally containing 1 further nitrogen atom wherein a -CH₂- group can optionally be replaced by a -C(O)-; which ring is optionally substituted on an available carbon or nitrogen atom by 1 substituent selected from hydroxy and R³;

R⁸ is selected from -OR⁵, (1-4C)alkyl, -C(O)(1-4C)alkyl, -C(O)NR⁴R⁵, (1-4C)alkylamino, di(1-4C)alkylamino, HET-3 wherein said ring is unsubstituted, (1-4C)alkoxy(1-4C)alkyl, hydroxy(1-4C)alkyl and -S(O)_pR⁵;

HET-4 is a 5- or 6-membered, C- or N- linked unsubstituted heteroaryl ring containing 1, 2, or 3 ring heteroatoms independently selected from O, N, and S;

p is independently at each occurrence 0, 1, or 2;

m is 1 and R² is in the para position relative to the ether linkage;

n is 0, 1, or 2.

Claim 23 (new): A compound of Formula (I) as claimed in Claim 22, or a salt, pro-drug, or solvate thereof, wherein HET-3 is a 4- to 6-membered ring.

Claim 24 (new): A compound of the Formula (I) as claimed in Claim 22, which is selected from:

- 3-[4-(azetidin-1-ylcarbonyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide;
- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-{4-[(4-methylpiperazin-1-yl)carbonyl]phenoxy}-N-1,3-thiazol-2-ylbenzamide;
- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-{4-[(4-methylpiperazin-1-yl)carbonyl]phenoxy}-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-[4-(morpholin-4-ylcarbonyl)phenoxy]benzamide;
- 3-[4-(azetidin-1-ylcarbonyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-[4-(pyrrolidin-1-ylcarbonyl)phenoxy]benzamide;
- 3-[4-(7-azabicyclo[2.2.1]hept-7-ylcarbonyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-({2-chloro-4-[(4-methylpiperazin-1-yl)sulfonyl]phenyl}oxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-({4-[(4-methylpiperazin-1-yl)sulfonyl]phenyl}oxy)-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-[4-(azetidin-1-ylcarbonyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(3-methyl-1,2,4-thiadiazol-5-yl)benzamide;
- 3-[4-(azetidin-1-ylcarbonyl)-2-chlorophenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-[4-(azetidin-1-ylcarbonyl)-2-fluorophenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-[4-(azetidin-1-ylcarbonyl)-2-(trifluoromethyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;


3-[4-(azetidin-1-ylcarbonyl)phenoxy]-5-[(1S)-2-methoxy-1-methylethoxy]-N-1H-pyrazol-3-ylbenzamide;
3-[4-(azetidin-1-ylcarbonyl)-2-chlorophenoxy]-5-[(1S)-2-methoxy-1-methylethoxy]-N-(5-methyl-1H-pyrazol-3-yl)benzamide;
3-[4-(azetidin-1-ylcarbonyl)-2-fluorophenoxy]-5-[(1S)-2-methoxy-1-methylethoxy]-N-(5-methyl-1H-pyrazol-3-yl)benzamide;
3-[4-(azetidin-1-ylcarbonyl)-2-fluorophenoxy]-5-[(1S)-2-methoxy-1-methylethoxy]-N-1H-pyrazol-3-ylbenzamide; and
3-[4-(azetidin-1-ylcarbonyl)phenoxy]-5-[(1S)-2-methoxy-1-methylethoxy]-N-(5-methyl-1H-pyrazol-3-yl)benzamide;
or a salt, pro-drug, or solvate thereof.

Claim 25 (new): A compound of Formula (I) as claimed in Claim 22, which is selected from:

3-[4-(azetidin-1-ylcarbonyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide;
3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-{4-[(4-methylpiperazin-1-yl)carbonyl]phenoxy}-N-1,3-thiazol-2-ylbenzamide;
3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-{4-[(4-methylpiperazin-1-yl)carbonyl]phenoxy}-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-[4-(morpholin-4-ylcarbonyl)phenoxy]benzamide;
3-[4-(azetidin-1-ylcarbonyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-[4-(pyrrolidin-1-ylcarbonyl)phenoxy]benzamide;
3-[4-(7-azabicyclo[2.2.1]hept-7-ylcarbonyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
3-({2-chloro-4-[(4-methylpiperazin-1-yl)sulfonyl]phenyl}oxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-({4-[(4-methylpiperazin-1-yl)sulfonyl]phenyl}oxy)-N-

(1-methyl-1H-pyrazol-3-yl)benzamide;
 3-[4-(azetidin-1-ylcarbonyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(3-methyl-1,2,4-thiadiazol-5-yl)benzamide;
 3-[4-(azetidin-1-ylcarbonyl)-2-chlorophenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
 3-[4-(azetidin-1-ylcarbonyl)-2-fluorophenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
 3-[4-(azetidin-1-ylcarbonyl)-2-(trifluoromethyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
 3-[4-(azetidin-1-ylcarbonyl)phenoxy]-5-[(1S)-2-methoxy-1-methylethoxy]-N-1H-pyrazol-3-ylbenzamide;
 3-[4-(azetidin-1-ylcarbonyl)-2-fluorophenoxy]-5-[(1S)-2-methoxy-1-methylethoxy]-N-(5-methyl-1H-pyrazol-3-yl)benzamide; and
 3-[4-(azetidin-1-ylcarbonyl)-2-fluorophenoxy]-5-[(1S)-2-methoxy-1-methylethoxy]-N-1H-pyrazol-3-ylbenzamide;
 or a salt, pro-drug, or solvate thereof.

Claim 26 (new): A compound of Formula (I), or a salt, pro-drug, or solvate thereof:



wherein:

R¹ is methoxymethyl;

R² is selected from -C(O)NR⁴¹R⁵¹, -SO₂NR⁴¹R⁵¹, and -S(O)_pR⁴¹;

HET-1 is a 5- or 6-membered, C-linked heteroaryl ring containing a nitrogen atom in the 2-position and optionally 1 or 2 further ring heteroatoms independently selected from O, N, and S; which ring is optionally substituted on an available carbon atom, or on a ring nitrogen

atom provided it is not thereby quaternised, with 1 or 2 substituents independently selected from R⁶;

HET-2 is a 4-, 5-, or 6-membered, C- or N-linked heterocyclyl ring containing 1, 2, 3, or 4 heteroatoms independently selected from O, N, and S, wherein a -CH₂- group can optionally be replaced by a -C(O)-, and wherein a sulphur atom in the heterocyclic ring may optionally be oxidised to a S(O) or S(O)₂ group, which ring is optionally substituted on an available carbon or nitrogen atom by 1 or 2 substituents independently selected from R⁷;

R³ is selected from halo, fluoromethyl, difluoromethyl, trifluoromethyl, methyl, methoxy, and cyano;

R⁴¹ is selected from (1-4C)alkyl substituted with 1 or 2 substituents independently selected from HET-2, -OR⁵, -SO₂R⁵, (3-6C)cycloalkyl (optionally substituted with 1 group selected from R⁷), and -C(O)NR⁵R⁵; (3-6C)cycloalkyl (optionally substituted with 1 group selected from R⁷); and HET-2;

R⁵¹ is hydrogen or (1-4C)alkyl;

R⁴ is selected from (1-4C)alkyl optionally substituted with 1 or 2 substituents independently selected from HET-2, -OR⁵, -SO₂R⁵, (3-6C)cycloalkyl (optionally substituted with 1 group selected from R⁷), and -C(O)NR⁵R⁵; (3-6C)cycloalkyl (optionally substituted with 1 group selected from R⁷); and HET-2;

R⁵ is hydrogen or (1-4C)alkyl;

or **R⁴** and **R⁵** together with the nitrogen atom to which they are attached may form a heterocyclyl ring system as defined by HET-3;

R⁶ is independently selected from (1-4C)alkyl, halo, hydroxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkylS(O)_p(1-4C)alkyl, amino(1-4C)alkyl, (1-4C)alkylamino(1-4C)alkyl, di(1-4C)alkylamino(1-4C)alkyl, and HET-4;

R⁷ is selected from -OR⁵, (1-4C)alkyl, -C(O)(1-4C)alkyl, -C(O)NR⁴R⁵, (1-4C)alkoxy(1-4C)alkyl, hydroxy(1-4C)alkyl, and -S(O)_pR⁵;

HET-3 is an N-linked, 4-, 5-, or 6-membered, saturated or partially unsaturated heterocyclyl ring, optionally containing 1 or 2 further heteroatoms independently selected from O, N, and S, wherein a -CH₂- group can optionally be replaced by a -C(O)- and wherein a sulphur atom in the ring may optionally be oxidised to a S(O) or S(O)₂ group; which ring is optionally substituted on an available carbon or nitrogen atom by 1 or 2 substituents independently

selected from R⁸; or

HET-3 is an N-linked, 7-membered, saturated or partially unsaturated heterocyclyl ring, optionally containing 1 further heteroatom independently selected from O, S, and N, wherein a -CH₂- group can optionally be replaced by a -C(O)- group and wherein a sulphur atom in the ring may optionally be oxidised to a S(O) or S(O)₂ group; which ring is optionally substituted on an available carbon or nitrogen atom by 1 or 2 substituents independently selected from R⁸; or

HET-3 is an N-linked, 6- to 10-membered bicyclic saturated or partially unsaturated heterocyclyl ring, optionally containing 1 further nitrogen atom wherein a -CH₂- group can optionally be replaced by a -C(O)-; which ring is optionally substituted on an available carbon or nitrogen atom by 1 substituent selected from hydroxy and R³;

R⁸ is selected from -OR⁵, (1-4C)alkyl, -C(O)(1-4C)alkyl, -C(O)NR⁴R⁵, (1-4C)alkylamino, di(1-4C)alkylamino, HET-3 (wherein said ring is unsubstituted), (1-4C)alkoxy(1-4C)alkyl, hydroxy(1-4C)alkyl, and -S(O)_pR⁵;

HET-4 is a 5- or 6-membered, C- or N- linked unsubstituted heteroaryl ring containing 1, 2, or 3 ring heteroatoms independently selected from O, N, and S;

p is independently at each occurrence 0, 1, or 2;

m is 1 and R² is in the para position relative to the ether linkage;

n is 0, 1, or 2.

Claim 27 (new): A compound of the formula (I) as claimed in Claim 26, which is selected from:

3-(4-[(2-methoxyethyl)amino]carbonyl}phenoxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide;

3-(4-[(1H-imidazol-2-ylmethyl)amino]carbonyl}phenoxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide;


3-(3-[(2-methoxyethyl)amino]carbonyl}phenoxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide;

3-(3-[(1H-imidazol-2-ylmethyl)amino]carbonyl}phenoxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide;

3-{[2-chloro-4-([2-(methyloxy)ethyl]amino)sulfonyl]phenyl}oxy}-5-[(1S)-2-methoxy-(1-

methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
 3-(4-[[[(2-methoxyethyl)amino]sulfonyl}phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide; and
 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-(4-[[[(1-methylpiperidin-4-yl)amino]carbonyl}phenoxy)-N-(3-methyl-1,2,4-thiadiazol-5-yl)benzamide;
 or a salt, pro-drug, or solvate thereof.

Claim 28 (new): A compound of the Formula (I), or a salt, pro-drug, or solvate thereof:



wherein:

R¹ is methoxymethyl;

R² is HET-2;

HET-1 is a 5- or 6-membered, C-linked heteroaryl ring containing a nitrogen atom in the 2-position and optionally 1 or 2 further ring heteroatoms independently selected from O, N, and S; which ring is optionally substituted on an available carbon atom, or on a ring nitrogen atom provided it is not thereby quaternised, with 1 or 2 substituents independently selected from R⁶;

HET-2 is a 4-, 5-, or 6-membered, C- or N-linked heterocyclyl ring containing 1, 2, 3, or 4 heteroatoms independently selected from O, N, and S, wherein a -CH₂- group can optionally be replaced by a -C(O)-, and wherein a sulphur atom in the heterocyclic ring may optionally be oxidised to an S(O) or S(O)₂ group, which ring is optionally substituted on an available carbon or nitrogen atom by 1 or 2 substituents independently selected from R⁷;

R³ is selected from halo, fluoromethyl, difluoromethyl, trifluoromethyl, methyl, methoxy, and cyano;

R⁴ is selected from hydrogen; (1-4C)alkyl optionally substituted with 1 or 2 substituents

independently selected from HET-2, $-OR^5$, $-SO_2R^5$, (3-6C)cycloalkyl (optionally substituted with 1 group selected from R^7), and $-C(O)NR^5R^5$; (3-6C)cycloalkyl (optionally substituted with 1 group selected from R^7); and HET-2;

R^5 is hydrogen or (1-4C)alkyl;

or R^4 and R^5 together with the nitrogen atom to which they are attached form a heterocyclyl ring system as defined by HET-3;

R^6 is independently selected from (1-4C)alkyl, halo, hydroxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkylS(O)_p(1-4C)alkyl, amino(1-4C)alkyl, (1-4C)alkylamino(1-4C)alkyl, di(1-4C)alkylamino(1-4C)alkyl, and HET-4;

R^7 is selected from $-OR^5$, (1-4C)alkyl, $-C(O)(1-4C)alkyl$, $-C(O)NR^4R^5$, (1-4C)alkoxy(1-4C)alkyl, hydroxy(1-4C)alkyl, and $-S(O)_pR^5$;

HET-3 is an N-linked, 4-, 5-, or 6-membered, saturated or partially unsaturated heterocyclyl ring, optionally containing 1 or 2 further heteroatoms independently selected from O, N, and S, wherein a $-CH_2-$ group can optionally be replaced by a $-C(O)-$ and wherein a sulphur atom in the ring may optionally be oxidised to an S(O) or S(O)₂ group; which ring is optionally substituted on an available carbon or nitrogen atom by 1 or 2 substituents independently selected from R^8 ; or

HET-3 is an N-linked, 7-membered, saturated or partially unsaturated heterocyclyl ring, optionally containing 1 further heteroatom independently selected from O, S, and N, wherein a $-CH_2-$ group can optionally be replaced by a $-C(O)-$ group and wherein a sulphur atom in the ring may optionally be oxidised to an S(O) or S(O)₂ group; which ring is optionally substituted on an available carbon or nitrogen atom by 1 or 2 substituents independently selected from R^8 ; or

HET-3 is an N-linked, 6- to 10-membered bicyclic saturated or partially unsaturated heterocyclyl ring, optionally containing 1 further nitrogen atom, wherein a $-CH_2-$ group can optionally be replaced by a $-C(O)-$; which ring is optionally substituted on an available carbon or nitrogen atom by 1 substituent selected from hydroxy and R^3 ;

R^8 is selected from $-OR^5$, (1-4C)alkyl, $-C(O)(1-4C)alkyl$, $-C(O)NR^4R^5$, (1-4C)alkylamino, di(1-4C)alkylamino, HET-3 wherein said ring is unsubstituted, (1-4C)alkoxy(1-4C)alkyl, hydroxy(1-4C)alkyl, and $-S(O)_pR^5$;

HET-4 is a 5- or 6-membered, C- or N- linked unsubstituted heteroaryl ring containing 1, 2, or 3

ring heteroatoms independently selected from O, N, and S;
p is independently at each occurrence 0, 1, or 2;
m is 1 and R² is in the para position relative to the ether linkage;
n is 0, 1, or 2.

Claim 29 (new): A compound of Formula (I), as claimed in Claim 28, which is:
3-[(1S)-2-methoxy-1-methylethoxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-[4-(1,2,4-oxadiazol-3-yl)phenoxy]benzamide;
or a salt, pro-drug, or solvate thereof.

Claim 30 (new): A compound of Formula (I) as claimed in Claim 18, Claim 22, Claim 26, or Claim 28 or a salt, pro-drug, or solvate thereof wherein R¹ has the (S) configuration.

Claim 31 (new): A compound of Formula (I) as claimed in Claim 18, Claim 22, Claim 26, or Claim 28 or a salt, pro-drug, or solvate thereof, wherein HET-1 is a 5-membered ring.


Claim 32 (new): A pharmaceutical composition comprising a compound as claimed in Claim 18, Claim 22, Claim 26, or Claim 28, or a salt, pro-drug, or solvate thereof, together with a pharmaceutically acceptable diluent or carrier.

Claim 33 (new): A method of treating GLK mediated diseases comprising administering an effective amount of a compound of Formula (I) as claimed in Claim 18, Claim 22, Claim 26, or Claim 28 or a salt, pro-drug, or solvate thereof, to a mammal in need of such treatment.

Claim 34 (new): The method of Claim 33, wherein the GLK mediated disease is type 2 diabetes.


Claim 35 (new): A process for the preparation of a compound of Formula (I) or a salt, pro-drug, or solvate thereof as claimed in Claim 18, Claim 22, Claim 26, or Claim 28, comprising:
(a) reacting an acid of Formula (III) or activated derivative thereof with a compound of Formula

(IV),



or


(b) reacting a compound of Formula (V) with a compound of Formula (VI),



wherein X^1 is a leaving group and X^2 is a hydroxyl group; or X^1 is a hydroxyl group and X^2 is a leaving group;


or

reacting a compound of Formula (V) with the intermediate ester of Formula (VII), wherein P^1 is a protecting group followed by ester hydrolysis and amide formation;



or

(c) reacting a compound of Formula (VIII) with a compound of Formula (IX)




(VIII)

(IX)

wherein X^3 is a leaving group or an organometallic reagent and X^4 is a hydroxyl group; or X^3 is a hydroxyl group and X^4 is a leaving group or an organometallic reagent;


or

reacting a compound of Formula (VIII) with the intermediate ester of Formula (X), followed by ester hydrolysis and amide formation;



or


(d) reacting a compound of Formula (XI) with a compound of Formula (XII),



wherein X^5 is a leaving group;

or

(e) when R^2 is of the formula $-C(O)NR^4R^5$, reacting a compound of the formula:



with a compound of the formula HNR^4R^5 ;

and thereafter, if necessary:

i) converting a compound of Formula (I) into another compound of Formula (I);

- ii) removing any protecting groups; and/or
- iii) forming a salt, pro-drug, or solvate.